

Biomarkers in Arthroplasty: A Systematic Review

Marty T. Mertens¹, Jasvinder A. Singh^{*,1,2,3,4}

¹Rheumatology Section, Medicine Service and ²Center for Chronic Disease Outcomes Research at the VA Medical Center, Minneapolis, MN, USA

³Division of Rheumatology, Department of Medicine, University of Minnesota, Minneapolis, MN, USA

⁴Departments of Health Sciences Research and Orthopedics, Mayo Clinic School of Medicine, Rochester, MN, USA

Abstract: We performed a systematic review of all MEDLINE-published studies of biomarkers in arthroplasty. Thirty studies met the inclusion criteria; majority evaluated biomarkers for osteolysis, aseptic prosthetic loosening, and prosthetic infections. Four studies reported an elevated Cross-linked N-telopeptides of type I collagen (urine or serum) in patients with osteolysis or aseptic prosthetic loosening when compared to appropriate controls. Two or more studies each found elevated C-reactive protein, erythrocyte sedimentation rate, and interleukin-6 in patients with infected prosthetic joints compared to controls. Most other biomarkers were either examined by single studies or had inconsistent or insignificant associations with outcomes. We conclude that the majority of the biomarkers currently lack the evidence to be considered as biomarkers for arthroplasty outcomes. Further studies are needed.

Keywords: Biomarkers, arthroplasty, systematic review.

INTRODUCTION

Total hip and knee arthroplasty (THA and TKA) represent a major advance in the treatment of refractory hip/knee joint pain. TKA and THA are performed very commonly and are projected to increase dramatically in numbers in the next two decades [1]. Increasing longevity, earlier age at arthroplasty, and excellent pain and functional outcomes with these procedures are cited as reasons for the increasing incidence of these procedures. A number of patients, however, have suboptimal outcomes. Complications may include refractory pain, peri-prosthesis osteolysis with aseptic loosening, heterotopic ossification, and infection, which may result in a failed arthroplasty. Evaluation for complications of arthroplasty are primarily clinical, with few prognostic or diagnostic tests easily available that allow early identification of this subgroup of patients with suboptimal outcomes.

Biomarkers are objective indicators of biologic processes, pathologic processes or pharmaceutical responses to therapeutic interventions [2]. A number of studies have assessed the prognostic and diagnostic utility of biomarkers in improving evaluation of outcomes following arthroplasty. The purpose of this systematic review is to summarize the data from the literature regarding biomarkers for arthroplasty outcomes. We specifically evaluated if there are any valid biomarkers for osteolysis or loosening of the prosthesis, prosthetic infections, pain/stiffness following arthroplasty, or other complications following arthroplasty.

METHODS

Study Design

All English language studies investigating biomarkers for various clinical outcomes of arthroplasty in humans were included in this review. Exclusion criteria included case reports, meta-analyses, and reviews. Studies that evaluated short term post-operative changes in biomarkers, biomarker changes related to medical therapies following arthroplasty, thromboembolic outcomes, and studies limited to pathology only were excluded as they were felt to be beyond the scope of this review.

Search Methods for Identification of Studies

A Cochrane librarian searched the MEDLINE electronic bibliographic database up to July 2008 using the MeSH terms “arthroplasty”, “replacement” and “biomarkers” and similar terms. The reference lists of identified publications were reviewed to identify any additional studies (Appendix 1 and 2).

Data Collection

Two reviewers (MM and JS) independently applied inclusion/exclusion criteria to all potential studies and extracted data and outcomes using a standardized form. Any disagreements were resolved by discussion between the two reviewers, referring to a third party if necessary. Review authors were not blinded to any features of studies, since unblinding has minimal impact on selection bias [3].

From each study, we extracted the following data – (1) study population, (2) methods and details of any interventions, (3) type of arthroplasty and implant, knee vs hip vs other, cemented vs noncemented, primary vs revision, and (4) types of biomarkers evaluated. Outcome data were

*Address correspondence to this author at the University of Alabama, Faculty Office Tower 805B, 510 20th Street S, Birmingham, AL 35294, USA; Tel: 205-934-8158; Fax: 205-966-9685;
E-mail: Jasvinder.md@gmail.com

Table 1. Summary of Biomarkers for Osteolysis

Outcome	Study (Hip, Knee, Both)	Osteolysis		No Osteolysis			Healthy Control		
		n	Mean (SD)	n	Mean (SD)	p-Value	n	Mean (SD)	p-Value
Collagen peptides									
Urinary cross-linked N-telopeptide of type I collagen	Antonioiu 2000 (hip)	11	NA	8 ^A	NA	p = 0.002	7	NA	p = 0.0297
	Von Schewelov 2006 (hip)	33	34 bone collagen equivalents (BCE)/nM creatinine (12)	127 ^A	29 BCE/nM Cr (15)	p = 0.06			
Serum C-terminal telopeptide of type I collagen	Arabmotagh 2006 (hip)	53	Significant correlation found between bone mineral density loss in ROI 7 at 1 year with week 3 CTX-1 values (p = 0.002)						
Chemokines, Cytokines & other									
Interleukin (IL)-1 β	Fiorio 2003 (hip)	8	2.15pg/ml (1.37)	10 ^A	2.26pg/ml (0.89)	p = 0.96	17	2.02pg/ml (1.25)	p = 0.79
	Hernigou 1999 (hip)	6	“Undetectable”	10 ^B	NA	NA	5	“undetectable”	NA
Tumor Necrosis Factor- α	Fiorio 2003 (hip)	8	4.32pg/ml (5.2)	10 ^A	3.84pg/ml (1.13)	p = 0.62	17	3.42pg/ml (2.12)	p = 0.95
	Hernigou 1999 (hip)	6	“undetectable”	10 ^B	NA	NA	5	“undetectable”	NA
IL-6	Fiorio 2003 (hip)	8	2.86pg/ml (1.95)	10 ^A	4.58pg/ml (4.02)	p = 0.24	17	3.88pg/ml (2.72)	p = 0.36
	Hernigou 1999 (hip)	6	41pg/ml (17)	10 ^B	22pg/ml (6.7)	p = 0.002	5	NA	NA
IL-11	Fiorio 2003 (hip)	8	0pg/ml (0)	10 ^A	1.22pg/ml (2.57)	p = 0.27	17	18.69pg/ml (10.55)	p = 0.001
transforming growth factor (TGF)- β	Fiorio 2003 (hip)	8	23175pg/ml (8774)	10 ^A	21120pg/ml (13657)	p = 0.50	17	23615pg/ml (10681)	p = 0.88
matrix metalloproteinase-1 (MMP-1)	Fiorio 2003 (hip)	8	3.69pg/ml (1.75)	10 ^A	4.10pg/ml (1.44)	p = 0.50	17	3.31pg/ml (1.7)	p = 0.52
prostaglandin E-2 (PGE-2)	Fiorio 2003 (hip)	8	1330pg/ml (1097)	10 ^A	2021pg/ml (1046)	p = 0.14	17	2893pg/ml (782)	p = 0.005
Bone Alkaline Phosphatase (ALP)	Arapmotagh 2006 (hip)	53	No significant correlation found between bone mineral density loss in ROI 7 at 1 year with bone ALP values						
Osteocalcin	Arapmotagh 2006 (hip)	53	No significant correlation found between bone mineral density loss in ROI 7 at 1 year with osteocalcin values						
C-reactive protein	Hernigou 1999 (hip)	6	10mg/L (2.4)	10 ^B	7.7mg/L (2.8)	p = 0.135	5	NA	NA

^AArthroplasty without osteolysis.^BArthroplasty without osteolysis, >10 years post-op.

ROI = region of interest.

extracted, where possible, as raw numbers, plus any summary measures where standard deviations or confidence intervals were given. If no raw results were available, specific outcomes were described in detail.

RESULTS

Eligible Studies

A total of 105 articles met initial criteria from the initial MEDLINE search completed in June 2007. The search was

updated in July 2008 and a total of 149 articles were identified (Fig. 1). A total of 30 studies met inclusion/exclusion criteria, 26 from the original search and four additional studies from the reference lists of these studies.

Osteolysis

A total of five studies evaluated biomarkers for osteolysis following knee or hip arthroplasty (summarized in Table 1). Most comparisons were made to populations with




Fig. (1). Summary of screening of eligible studies.

arthroplasty with no osteolysis and to healthy controls without arthroplasty.

Collagen telopeptides: Three studies evaluated collagen telopeptides as biomarkers for osteolysis following hip arthroplasty. Antoniou *et al.* [4] reported significantly higher urinary cross-linked N-telopeptides of type-T collagen in patients with osteolysis following hip arthropathy compared to controls. von Schewelov *et al.* [5] found a near significant increase in urinary cross-linked N-telopeptide in the osteolysis versus non-osteolysis group ($p=0.06$), which became significant after adjusting for age, gender, and time after surgery. Arabmotlagh *et al.* [6] found serum C-terminal

telopeptides of type I collagen 3 weeks post-operatively to be significantly correlated with bone loss one year after hip arthroplasty.

Interleukin-1 β (IL-1 β) and Tumor Necrosis Factor-alpha (TNF- α): Neither of the two studies that evaluated serum IL-1 β or TNF- α as a biomarker for osteolysis following hip arthroplasty found a significant difference between patients with osteolysis compared to controls [7, 8].

IL-6: Two studies [7, 8] evaluated IL-6 as a biomarker for osteolysis following hip arthroplasty. Hernigou *et al.* [8]

Table 2. Summary of Biomarkers for Loosening of Prosthesis

Outcome	Study (Hip, Knee, Both)	Loosening		Non-Loosened ^A		p-Value	Healthy Control		p-Value
		n	Mean (SD)	n	Mean (SD)		n	Mean (SD)	
Collagen peptides									
Serum C-terminal propeptide of type I procollagen	Schneider 1998 (hip)	50	107.5 ng/ml [#] (70.4)	50	82.2 ng/ml [#] (32.8)	NS			
	Savarino 2005 (hip)	21	98.40 ng/ml (57.47)	18	124.73 ng/ml (53.12)	p = 0.04	30	172.81 ng/ml (59.7)	NA
	Li 2004 (knee)	15 ^C		25 ^D	No statistical difference between unstable and stable joints from 3 – 24 months post-op (p = 0.79, ANOVA)				
Serum N-terminal propeptide of type I procollagen	Wilkinson 2003 (hip)	17 ¹ 18 ²	46.9 ug/ml ¹ (NA) 45.6 ug/ml ²	32 ¹ , 31 ²	37.5 ug/ml ¹ (NA) 38.0 ug/ml ² (NA)	NS			
Serum cross-linked N-terminal telopeptide	Savarino 2005 (hip)	21	22.9nM bone collagen equivalents (BCE)/L [*] (6.05)	18	17.92nMBCE/L [*] (4.24)	p = 0.02	30	17.99nMBCE/L [*] (3.61)	NA
Serum cross-linked C-terminal telopeptide	Li 2004 (knee)	15 ^C		25 ^D	Statistically higher levels in unstable joints vs stable joints from 6 – 24 months post-op (p = 0.02, ANOVA)				
	Wilksinson 2003 (hip)	17 ¹ 18 ²	0.43 ng/ml ¹ (NA) 0.45 ng/ml ² (NA)	32 ¹ , 31 ²	0.33 ng/ml ¹ (NA) 0.33 ng/ml ² (NA)	NS			
Serum Type III procollagen peptide	Moreschini 1997 (hip)	9	NA	13	NA	NS			
Urine cross-linked N-telopeptide of type I collagen	Schneider 1998 (hip)	50	NA	50	NA	p <0.0001			
	Wilkinson 2003 (hip)	17 ¹ 18 ²	61.0nM BCE/mM Cr ^{1,*} (NA) 62.3nM BCE/mM Cr ^{2,*} (NA)	32 ¹ , 31 ²	39.9 nM BCE/mM Cr ^{1,*} (NA) 42.8nM BCE/mM Cr ^{2,*} (NA)	p = 0.02 ¹ NS ²			
Chemokines, Cytokines & other									
Tumor Necrosis Factor- α	Granchi 1998 (hip)	35	0.06 pg/ml (0.35)				25	0 (0)	NS
	Hundri-Haspl 2006 (both)	50	32.7 pg/ml (32.4)	50	22.9 pg/ml (18.7)	p <0.01	50	3.9 pg/ml (3.9)	p <0.001
Interleukin (IL)-1 β	Granchi 1998 (hip)	35	0.33 pg/ml (0.59)				25	0.37 pg/ml (0.5)	NS

(Table 2) contd.....

Outcome	Study (Hip, Knee, Both)	Loosening		Non-Loosened ^A		p-Value	Healthy Control		p-Value
		n	Mean (SD)	n	Mean (SD)		n	Mean (SD)	
	Hundri-Haspl 2006 (both)	50	3.7 pg/ml (5.5)	50	1.5 pg/ml (2)	p <0.001	50	0.7 pg/ml (1.1)	p <0.001
	Moreschini 1997 (hip)	9	NA (detectable in 4/9 pts)	13	NA (detectable in 1/13 pts)	NS			
IL-6	Granchi 1998 (hip)	35	28.9 pg/ml (65)				25	35.3 (45)	NS
	Streich 2003	23	3.97 pg/ml (5.33)	23	4.08 pg/ml (6.11)	NS			
IL-8	Hundri-Haspl 2006 (both)	50	14.7 pg/ml (9)	50	8.1 pg/ml (4.7)	p <0.001	50	4.2 pg/ml (1.3)	p <0.001
IL-2 receptor	Granchi 1998 (hip)	35	86.6 pg/ml (35.5)				25	83.9 pg/ml (35)	NS
	Streich 2003	23	469.2U/ml (155.2)	23	515U/ml (160.5)	NS			
Granulocyte-macrophage colony stimulating factor (GM-CSF)	Granchi 1998 (hip)	35	29.6 pg/ml (41.4)				25	4.4 pg/ml (10)	NS
	Streich 2003	23	0.21 pg/ml (1.0)	23	not detectable	NS			
Osteocalcin	Wilkinson 2003 (hip)	17 ¹ 18 ²	22.2 ng/ml ¹ (NA) 20.7 ng/ml ² (NA)	32 ¹ 31 ²	19.0 ng/ml ¹ (NA) 20.6 ng/ml ² (NA)	NS			
	Schneider 1998 (hip)	50	NA	50	NA	p <0.02			
	Li 2004 (knee)	15 ^C		25 ^D	Statistically higher levels in unstable joints vs stable joints at 12 months post-op only (p = 0.03)				
Elastase	Streich 2003	23	58.9 ng/ml (46.8)	23	56.6 ng/ml (45.0)	NS			
Hyaluronic acid	Moreschini 1997 (hip)	9	779.3ug/L (475.8)	13	112.9ug/L (42.5)	p <0.05			
Bone Alkaline Phosphatase	Wilkinson 2003 (hip)	17 ¹ 18 ²	32.6 U/L ¹ (NA) 28.5 U/L ² (NA)	32 ¹ 31 ²	27.9 U/L ¹ (NA) 28.2 U/L ² (NA)	NS			
	Schneider 1998 (hip)	50	123.8 U/L [#] (42.5)	50	110.4 U/L [#] (28.0)	NS			
Nuclear factor-kB ligand (RANKL)	Granchi 2006 (hip)	36	1483 pg/ml (1179)	33	3312 pg/ml (2211)	p <0.002	20	1606 pg/ml (1245)	NS
Osteoprotegerin	Granchi 2006 (hip)	36	4198 pg/ml (286)	33	2397 pg/ml (1632)	p <0.0001	20	2586 pg/ml (1082)	<0.0001

(Table 2) contd.....

Outcome	Study (Hip, Knee, Both)	Loosening		Non-Loosened ^A		p-Value	Healthy Control		p-Value
		n	Mean (SD)	n	Mean (SD)		n	Mean (SD)	
Urine free deoxypyridinoline	Wilkinson 2003 (hip)	17 ¹ 18 ²	8.2nmol/mM Cr ¹ (NA) 8.4nmol/mM Cr ² (NA)	32 ¹ , 31 ²	7.6nmol/mM Cr ¹ (NA) 7.5nmol/mM Cr ² (NA)	NS			
Urine deoxypyridinoline	Schneider 1998 (hip)	50	NA	50	NA	p<0.001			
urine pyridinoline	Schneider 1998 (hip)	50	NA	50	NA	p<0.001			
Synovial Analyses:									
Osteocalcin	Synpniewska 2002 (hip)	8	4.2 ng/ml [#] (NA)	29 ^B	22.6 ng/ml [#] (NA)	p = 0.01			
B-crosslaps	Synpniewska 2002 (hip)	10	0.8 ng/ml [#] (NA)	36 ^B	1.3 ng/ml [#] (NA)	p = 0.004			
Interleukin (IL)-1 β	Kovacik 2000 (knee)	20	6.25 pg/ml (4.34)	7 ^B	0.25 pg/ml (0.01)	p <0.0001			
	Nibrant 1999 (hip)	26	17 pg/ml [#] (NA)	12	7 pg/ml [#] (NA)	0.18			
IL-1 α	Synpniewska 2002 (hip)	8	1.9 pg/ml [#] (NA)	10 ^B	1.4 pg/ml [#] (NA)	NS			
IL-6	Nibrant 1999 (hip)	27	652 pg/ml [#] (NA)	1	1800 pg/ml [#] (NA)	NA			
IL-8	Synpniewska 2002 (hip)	9	870 pg/ml [#] (NA)	26 ^B	340 pg/ml [#] (NA)	p = 0.01			
IL-10	Synpniewska 2002 (hip)	9	14.3 pg/ml [#] (NA)	24 ^B	4.0 pg/ml [#] (NA)	p = 0.004			
Tumor Necrosis Factor- α	Nibrant 1999 (hip)	38	31 pg/ml [#] (NA)	12	25 pg/ml [#] (NA)	0.22			
Bone Alkaline Phosphatase	Synpniewska 2002 (hip)	7	68.7U/L [#] (NA)	20 ^B	98.1U/L [#] (NA)	p = 0.01			
TNF-related activation protein (TRAP)	Kovacik 2000 (knee)	20	51.12U/L (25.75)	7 ^B	14.07U/L (3.90)	p <0.0001			
Chitinase 3-like protein 1 (YKL-40)	Kawasaki 2001 (hip)	5	3265 ng/ml [#] (NA)	19 ^B	886 ng/ml [#] (NA)	p = 0.015			

^ANanomoles of bone collagen equivalents. ^AArthroplasty without evidence of osteolysis or loosening, unless otherwise noted. ^BOA patients, undergoing primary arthroplasty.

^CPotentially unstable knees, determined by MTPM >0.2mm on radiostereometric analysis (RSA); ^DStable knees, determined by MTPM <0.2mm on RSA. ¹Femoral implant loosening. ²Pelvic implant loosening. [#]Median values (means not provided). MTPM = the largest three-dimensional translation of the tibial component.

reported significantly higher IL-6 levels in patients with osteolysis compared to controls, while, in contrast, Fiorto *et al.* [7] did not find a significant difference in IL-6 levels with similar comparison groups.

Other: Biomarkers evaluated by single studies included bone alkaline phosphatase (bone ALP) and osteocalcin [6]; IL-11, transforming growth factor-beta (TGF- β), matrix metalloproteinase-1 (MMP-1) and prostaglandin E-2 (PGE-2) [7]; and C-reactive protein (CRP) [8]. Only IL-11 was significantly lower in patients with osteolysis when compared to healthy controls and those without osteolysis.

Loosening

A total of 14 studies evaluated biomarkers for loosening of the prosthesis following knee or hip arthroplasty compared to healthy controls or patients with non-loosened prostheses (summarized in Table 2). Nine studies evaluated serum biomarkers and five, synovial fluid biomarkers.

Serum Biomarkers

Collagen Telopeptides: Five studies evaluated various collagen peptides as a biomarker for prosthetic loosening. Serum levels of a C-terminal propeptide of type I procollagen (PICP) were not significantly different between clinically loosened hip prosthetic joints and controls [9] or between unfixed vs fixed knee prostheses [10]. However, Savarino *et al.* [11] demonstrated a significantly lower PICP concentrations in loosened hip prostheses compared to stable hip implants. Serum N-terminal propeptide of type I procollagen (PINP) levels did not differ between patients with hip prostheses loosening and controls [12].

Savarino *et al.* [11] reported significantly higher serum cross-linked N-terminal telopeptide (NTx) levels in loosened hip prostheses compared to controls. Li *et al.* [10] found significantly higher levels of serum cross-linked C-terminal telopeptides (ICTP) in unfixed knee arthroplasties when compared to fixed joints. Conversely, Wilkinson *et al.* [12] found no difference in ICTP levels between patients with loosened hip prostheses and controls. Moreschini *et al.* [13] found no significant difference in type III procollagen peptide levels in patients with loosened hip prostheses when compared to controls with osteoarthritis.

Schneider *et al.* [9] and Wilkinson *et al.* [12] both found significantly increased levels of urinary crosslinked N-telopeptides of type I collagen in patients with loosened hip arthroplasty when compared to stable prostheses.

TNF- α : Two studies [14, 15] evaluated TNF- α as a biomarker for loosening of arthroplasty. Hundri-Haspl *et al.* [15] noted significant increase in TNF- α values in patients with loosened knee and hip arthroplasty when compared to controls, while Granchi *et al.* [14] found no difference in TNF- α levels in similar comparison groups.

IL-1 β : Three studies [13-15] evaluated serum IL-1 β as a biomarker for arthroplasty loosening. Hundri-Haspl *et al.* [15] noted significant increase in IL-1 β in patients with loosened knee and hip arthroplasty when compared to controls. Granchi *et al.* [14] did not find a difference in IL-1 β when comparing loosened hip arthroplasty to healthy controls. However, on subgroup analysis, they found a

significant increase in IL-1 β in patients with loosened titaniumaluminium-vanadium (TiAlV) cemented hip prostheses in comparison to healthy controls as well as loosened cobalt-chromium (CrCoMo) hip implants (cemented and uncemented) and uncemented TiAlV implants. Moreschini *et al.* [13] reported detectable IL-1 β in 4 of 9 patients with loosened hip implants compared to only 1 of 13 patients with fixed hip implant and 0 of 13 patients with OA prior to arthroplasty, though no statistical analysis was performed.

IL-6: Two studies [14, 16] found no significant differences in IL-6 levels in patients with loosened hip arthroplasty compared to controls.

Granulocyte-macrophage colony stimulating factor (GM-CSF): Streich *et al.* [16] demonstrated detectable levels of GM-CSF in only 1 of 23 patients with loosened hip arthroplasty and in 0 of 23 of fixed hip implant patients. Granchi *et al.* [14] reported significantly higher levels of GM-CSF in patients with a loosened hip arthroplasty when compared to healthy controls. This significant increase was demonstrated in all subgroups (cemented/uncemented TiAlV implants, cemented CrCoMo implants) except loosened uncemented CrCoMo implants which were similar to controls.

Osteocalcin: Schneider *et al.* [9] reported significantly higher osteocalcin levels in loosened versus stable prostheses, while Wilkinson *et al.* [12] found no difference with the same comparison. Li *et al.* [10] found significantly higher osteocalcin levels in patients with potentially unstable knee prostheses compared to patients with stable implants at 12 months, but no differences were noted at 24 months.

Bone alkaline phosphatase (ALP): Bone-specific ALP levels were similar in patients with and without loosening of hip prosthesis in two studies [9, 12].

Urine deoxypyridinoline: Schneider *et al.* [9] found significantly higher urine deoxypyridinoline levels in patients with loosened hip implants compared to stable implants, while Wilkinson *et al.* [12] reported no significant difference in similar comparison groups.

IL-2 receptor: Two studies [14, 16] found no significant differences in IL-2 receptor concentrations in patients with prosthesis loosening compared to controls.

Other: The evidence for each of the following was based on single studies. Significantly higher levels of osteoprotegerin (OPG) and RANKL [17], hyaluronic acid [13], urine pyridinoline [9] and IL-8 [15] were found in patients with loosened hip implants compared to their respective controls. Elastase levels were not significantly different between patients with loosened implant and controls [16].

Synovial Biomarkers

Four studies evaluated various biomarkers in synovial fluid as a marker for loosening of the prosthetic joint when compared to controls.

IL-1 β : Nivbrant *et al.* [18] found significantly higher synovial IL-1 β in loosened hip implants compared to osteoarthritis patients, but were similar to patients with fixed

Table 3. Summary of Biomarkers for Infected Prostheses

Outcome	Study (Hip, Knee, Both)	Infected Prosthesis		Control Group		p-Value
		n	Mean (SD)	n	Mean (SD)	
Interleukin (IL)-6	Bottner 2007 (both)	21	48.2 pg/ml (58.9)	57 ^A	10.6 pg/ml (24.2)	p = 0.0001
	Di Cesare 2005 (both)	17	50.3 pg/ml (40.9)	41 ^A	3.2 pg/ml (3.6)	p <0.01
Tumor Necrosis Factor- α (TNF- α)	Bottner 2007 (both)	21	39.7 ng/ml (23.4)	57 ^A	23.5 ng/ml (13.3)	p = 0.0011
Procalcitonin	Bottner 2007 (both)	21	1.5 ng/ml (3.3)	57 ^A	0.1 ng/ml (0.4)	p = 0.0033
C-reactive protein (CRP)	Bottner 2007 (both)	21	12.0mg/dl (9.0)	57 ^A	1.1mg/dl (1.5)	p <0.0001
	Di Cesare 2005 (both)	17	142.0mg/L (145.3)	41 ^A	6.4mg/L (7.4)	p <0.01
	Nilsdotter 2007 (hip)	22	24mg/L [#] (NA)	56 ^B	<10mg/L [#] (NA)	p <0.001
	Spangehl 1999 (hip)	prospective analysis of 142 revision hip arthroplasties: CRP >10mg/L had sensitivity of 0.96 and specificity of 0.92 (specific data not provided)				
Erythrocyte sedimentation rate (ESR)	Bottner 2007 (both)	21	63.8mm (24.4)	57 ^A	19.9mm (18.9)	p <0.0001
	Di Cesare 2005 (both)	17	82.0mm (33.1)	41 ^A	30.4mm (25.9)	p <0.01
	Nilsdotter 2007 (hip)	22	36mm [#] (NA)	55 ^B	14mm [#] (NA)	p <0.001
	Spangehl 1999 (hip)	prospective analysis of 171 revision hip arthroplasties: ESR >30mm had sensitivity of 0.82 and specificity of 0.85 (specific data not provided)				
Bacterial Markers:						
IgG lipoteichoic acid	Rafiq 2000 (both)	15 ^D	64,000 titer (47,451)	32 ^C	8469 titer (5136)	p <0.0001
IgM lipoteichoic acid	Rafiq 2000 (both)	15 ^D	3992 titer (4094)	32 ^C	3533 titer (3136)	p = 0.79
ica A operon	Frank 2004	44 ^E	20/44 isolates (46%) with ica A	23 ^F	7/23 isolates (30%) with ica A	p = 0.23
	Frank 2004	31 ^G	17/31 isolates (55%) with ica A	9 ^H	5/9 isolates (56%) with ica A	p = 1.0
	Galdbert 2000 (both)	54 ^G	44/54 isolates (81.5%) with ica A	23 ^I	4/23 isolates (17.4%) with ica A	p = 0.01
Biofilm production	Galdbert 2000 (both)	54 ^G	34/54 isolates (62.9%) with biofilm production	23 ^I	2/23 isolates (8.6%) with biofilm production	
Other operons	Galdbert 2000 (both)	54 ^G		23 ^I	no significant difference in frequency of the following operons: clfA, atlE, fbe, finbA, cna	
Synovial markers:						
Cytokines (TNF- α , IL-6, IL-1 β)	Nilsdotter 2007 (hip)	17		47 ^B	significantly higher synovial levels in infected group vs aseptic loosening group (data not provided)	
Lactate	Nilsdotter 2007 (hip)	17	5.8mmol/L [#] (NA)	45 ^B	5.4mmol/L [#] (NA)	p = 0.40

^AAseptic revision arthroplasties, ^BAseptic loosening of hip arthroplasty, ^CRecent closed fracture or osteoarthritis undergoing primary osteoarthritis of the hip, ^DInfections with gram positive bacteria, ^ECoagulase negative staph cultures from prosthetic infections, ^FCoagulase negative staph cultures from non-infected prostheses, ^GStaph epidermidis cultures from prosthetic infections, ^HStaph epidermidis cultures from non-infected prostheses, ^IStaph epidermidis cultures from skin flora on healthy individuals, [#]Median values (mean values not provided)

mein CRP 3.44 mg/L für Hüft TEP ok, insbes. weil er schon lange bei diesem Wert ist, sagt Koch 15.5.2015.

hip prostheses. Similarly, Kovacik *et al.* [19] reported significantly higher synovial IL-1 β levels in patients with loosened knee prostheses compared to osteoarthritis patients.

Other: Nivbrant *et al.* [18] demonstrated significantly higher synovial TNF- α levels in loosened prosthetic hip joints when compared to OA patients, though not when compared to fixed hip implants. Synovial IL-6 levels were similar between all of these groups, though too few observations were available for comparisons. Kawasaki *et al.* [20] noted significantly higher YKL-40 levels, a growth factor for chondrocytes and fibroblasts, in failed hip implants when compared to OA patients, though these levels were similar to patients with osteonecrosis of the hip. Other biomarkers found to be significantly elevated in loosened prostheses when compared to OA patients in single studies were TNF-related activation protein (TRAP) concentrations [19], IL-8 [21], and IL-10 [21]. Lower concentrations of osteocalcin, B-crosslaps, and bone ALP were also noted in loosened hip prostheses when compared to OA patients in a single study [21]. IL-1 α concentrations were similar in both groups [21].

Prosthetic Infections

A total of seven studies evaluated biomarkers for infection of the prosthesis following knee or hip arthroplasty (summarized in Table 3). Three studies, Rafiq *et al.* [22], Galdbert *et al.* [23], and Frank *et al.* [24] evaluated markers for specific infections of prosthetic joints and four studies evaluated diagnostic biomarkers for prosthetic infections.

C-reactive protein (CRP) and Erythrocyte sedimentation rate (ESR): Three separate studies [25-27] each found significantly higher CRP and ESR levels in patients infected prosthetic joints compared to aseptic revisions. Spangehl *et al.* [28] prospectively analyzed patients undergoing hip revisions and noted a sensitivity of 0.96 and specificity of 0.92 for CRP levels >10 mg/L for infected versus aseptic revisions. For ESR>30, the respective sensitivity was 0.82 and specificity, 0.85.

IL-6: Two studies [25, 26] each noted significantly higher levels of serum IL-6 in infected prosthetic joints when compared to aseptic joints undergoing revision arthroplasty.

Other cytokines: Bottner *et al.* [26] noted elevated levels of serum TNF- α and procyclitin in patients with infected prosthetic joints compared to aseptic prosthetic revisions.

Synovial fluid markers: Nilsdotter *et al.* [27] noted significantly higher synovial fluid concentrations of TNF- α , IL-1 β , and IL-6 in those with infected prosthetic hip joints when compared to aseptic controls.

Bacterial markers: Galdbert *et al.* [23] evaluated bacterial genes from *Staphylococcus epidermidis* strains cultured from prosthetic infections as compared to *S. epidermidis* strains from skin flora of healthy controls. They found significantly greater frequency of biofilm production and the ica A operon in infected prostheses. In contrast, Frank *et al.* [24] found no difference in frequency of the ica A operon in coagulase-

negative staph species cultured from prosthetic infections in comparison to strains obtained from prosthetic joints without evidence of infection.

Rafiq *et al.* [22] reported significantly elevated titers of IgG to cellular lipoteichoic acid in patients with prosthetic joint infections as compared to patients with osteoarthritis or closed fractures. IgM titers were similar between these groups.

OTHER OUTCOMES

Pain/Stiffness

Hall *et al.* [29] found a significant correlation between peak CRP levels post-operatively and pain at time of discharge as well as between peak IL-6 levels post-operatively and time to be able to ambulate 10 and 25m following hip arthroplasty. No significant correlations were found between CRP or IL-6 levels post-operatively to 1 and 6 month Western Ontario and McMaster universities osteoarthritis index (WOMAC) values, a measure of arthritis-related disability.

Wozniak *et al.* [30] prospectively found significantly higher nitrous oxide (NO) production from stimulated peripheral neutrophils 2 months and 2-3 years post-operatively in patients with pain compared to patients without pain following primary knee or hip arthroplasty.

Heterotopic Ossification

Wilkinson *et al.* [31] noted significantly greater levels of C-telopeptide of type-I collagen, osteocalcin, and N-terminal propeptide of type-I procollagen in patients who developed heterotopic ossification at 26 weeks when compared to those without ossification. No differences were noted in bone ALP, urinary cross-linked N-telopeptide of type-I collagen, and urinary free deoxypyridinoline.

Polyethylene Wear

Messieh *et al.* [32] noted a significant correlation between synovial lactate dehydrogenase (LDH) levels and polyethylene wear in eleven patients undergoing revision knee arthroplasty.

Hospital Outcomes

Ackland *et al.* [33] prospectively noted patients with a preoperative CRP >3mg/L had a significantly longer mean hospital stay as compared to those with CRP <3 mg/L (7.5 vs 6.0 days, respectively) following knee or hip arthroplasty.

DISCUSSION

This systematic review of biomarkers in arthroplasty summarized the findings from 30 studies of clinical outcomes following knee and/or hip arthroplasty. Despite the enormous success rates of knee and hip arthroplasties, patients may face a number of potential complications post-operatively and during long-term follow-up, including periprosthetic osteolysis, implant loosening, and joint infections. Diagnostic and prognostic biomarkers have the potential to

provide an early, accurate, and noninvasive diagnosis of these undesired outcomes as well as help design interventions to prevent some of these complications.

The most frequently studied arthroplasty outcomes with regards to biomarkers were osteolysis and prosthesis loosening. Osteolysis and subsequent aseptic joint loosening is the most common reason for joint revision in hip arthroplasty in the United States.

In our review of the published literature, only cross-linked Ntelopeptides of type I collagen (NTX-1) (serum or urine) was significantly associated with loosening/osteolysis, in more than two studies with appropriate controls (without loosening/osteolysis and/or healthy controls) [4, 9, 11, 12]. Von Schewelov *et al.* found an insignificant difference between NTX-1 levels in patients with presence of osteolysis when compared to controls ($p = 0.06$), though became significant after adjustment for age, gender, and time after surgery.

Serum cytokines, such as serum IL-1 β or TNF- α , that have been suspected to be associated with development of osteolysis and loosening [35-38] had inconsistent associations with loosened prostheses [14, 15] and no association with osteolysis [7, 8]. We postulate that this may be due to lack of well-designed studies rather than lack of association. A significant immunological response occurs in the periprosthetic tissue in many patients with prosthetic loosening, and it has been thought that this inflammatory process may contribute to the development of osteolysis and aseptic loosening [35-38]. Other biomarkers for osteolysis or prosthesis loosening were either studied by single studies or had inconsistent or insignificant differences.

Prosthetic infections lead to significant morbidity, are difficult to diagnose, and may lead to significant disability. CRP and ESR, both of which are currently used clinically in the diagnosis and follow-up of patients with prosthetic joint infections, were consistently found to be significantly elevated in septic prosthetic joints as compared to aseptic joint revision controls in a number of studies [25-28]. Interleukin-6 was consistently found to be elevated in septic prostheses as compared to aseptic revision controls [25, 26].

A number of other arthroplasty outcomes included in this review, such as heterotopic ossification, pain following arthroplasty, post-operative hospital outcomes, and

polyethylene wear, were evaluated only by one or two studies. Because of this, it is difficult to make adequate assessments for their potential use as biomarkers for these outcomes.

This review had a number of limitations, mostly related to limitations of the included studies. The large variability of the biomarkers measured in the included studies with only a few studies replicating the results for each biomarker in more than one study/population, limited our ability to draw conclusions for most biomarkers. The majority of the included studies were case control studies with small sample sizes, with half of the included studies containing <50 patients. A number of these studies measured multiple biomarkers at multiple outcomes, making them prone to type I error, i.e., finding a positive result just by chance. Well-designed studies of larger, welldefined cohorts are needed to improve our understanding of the utility of these biomarkers in predicting arthroplasty outcomes.

In conclusion, this review evaluated 30 studies that measured different biomarkers for knee and hip arthroplasty outcomes. While a few biomarkers – namely cross-linked Ntelopeptides of type I collagen for prosthesis loosening or osteolysis and CRP, ESR, and IL-6 for infected prostheses – were found consistently significantly different from control groups in the included studies, most biomarkers lack the evidence to be considered adequate markers for the arthroplasty outcomes at the present time. However, this is an exciting area of research that has the potential to impact and improve arthroplasty outcomes. Further studies are necessary, likely with larger study populations and longer prospective follow-up, to determine which of these biomarkers will have clinical application in diagnosis and/or prognosis of arthroplasty outcomes.

ACKNOWLEDGEMENTS

We thank Indy Rutks of the Cochrane Library group at the Minneapolis Veterans Affairs Medical center, Minneapolis for performing and updating the search.

GRANT SUPPORT

Supported by the NIH CTSA Award 1 KL2 RR024151-01 (Mayo Clinic Center for Clinical and Translational Research).

Appendix 1. Search strategy for Systematic Review Performed in 2007 (Updated in July 2008)

Database: Ovid MEDLINE(R) <1950 to July Week 4 2007>

Search Strategy:

- 1 Biological Markers/ or biomarkers.mp. (79469)
- 2 exp Arthroplasty, Replacement, Knee/ or exp Arthroplasty, Replacement/ or exp Arthroplasty, Replacement, Hip/ or exp Arthroplasty/ or exp Arthroplasty, Replacement, Finger/ or exp Joint Prosthesis/ or arthroplasty.mp. (39802)
- 3 1 and 2 (125)
- 4 limit 3 to english language (105)

Appendix 2. Summary of the Included Studies

Study	Study Design/ Level of Evidence	Comparisons	Outcomes	Other Notes
Antonioiu 2000	case control study/Level III	Hip osteolysis (n = 11), hip arthroplasty without osteolysis (n = 8), healthy controls (n = 7), hip osteolysis following alendronate therapy x6 weeks (n = 10)	Urinary cross-linked N-telopeptide of type I collagen;	Longer average time from original operation in osteolysis group (8.8yrs) vs. fixed implants (2.4yrs)
Von Schewelov 2006	case control study/Level III	Hip osteolysis (n = 33); hip arthroplasty without osteolysis (n = 127)	urinary cross-linked N-telopeptide of type I collagen	Post-operative time in osteolysis was 73 months vs. 42 months in non-osteolysis group.
Arabmotagh 2006	Prospective cohort study/Level II	53 patients with unilateral THA for OA; followed with bone density scans on 2, 4, 6, and 12mos post-op and correlated to marker measured at 3, 6, 16, and 24 weeks post-op.	Serum C-terminal telopeptide of type I collagen; bone ALP, osteocalcin	Numerous comparisons from BMD and biomarkers lessen reliability of findings.
Hernigou 1999	case control study/Level III	Hip osteolysis, >10yrs postop (n = 6); hip arthroplasty without osteolysis, >10yrs post-op (n = 10); hip arthroplasty without osteolysis, <6yrs post-op (n = 6); healthy controls (n = 5)	IL-B, TNF-a, IL-6, CRP	
Fiorito 2003	case control study/Level III	Hip osteolysis (n = 8); hip arthroplasty without osteolysis (n = 10); healthy controls (n = 17)	IL-1B, IL-6, TNF-a, TGF-B, PGE-2, MMP-1, IL-11	
Schneider 1998	case control study/Level III	Aseptic loosening of hip implant (n = 50); fixed hip implant (n = 50)	Bone ALP, osteocalcin, serum C-terminal propeptide of type I procollagen, urine cross-linked N-telopeptide of type I collagen, urine pyridinoline, urine deoxypyridinoline	
Savarino 2005	case control study/Level III	Aseptic loosening of hip implant (n = 21), fixed hip implant (n = 18), OA patients at time of primary arthroplasty (n = 17), healthy controls (n = 30)	Serum cross-linked N-telopeptide of type I collagen, serum C-terminal propeptide of type I procollagen	Healthy controls not matched for age or gender and not used in statistical comparisons
Li 2004	prospective cohort study/Level II	45 patients followed prospectively after TKA; biomarker measured at 1 week and 3, 6, 12, and 24 months post-op; compared to quality of fixation measured by RSA ¹ – MTPM ² >0.2mm represented unstable fixation, <0.2mm represented stable fixation	serum C-terminal propeptide of type I procollagen; serum cross-linked C-terminal telopeptide of type I collagen; osteocalcin	A part of an 80 patient randomized prospective study comparing cemented to uncemented fixation of tibial component for TKA, 45 agreed for blood tests; excluded patients without RSA data at 1 or 2 years (5 patients)
Wilkinson 2003	case control study/Level III	Aseptic loosening of hip implant (n = 26); stable fixation of hip implant (n = 23)	Bone ALP, serum cross-linked C-terminal telopeptide, osteocalcin, serum N-terminal propeptide of type I procollagen, urine cross-linked N-telopeptide of type I collagen, free deoxypyridinoline	Only provided data for either femoral or pelvic loosening when compared to fixed implants
Moreschini 1997	case control study/Level III	Aseptic loosened hip prosthesis (n = 9); fixed hip prosthesis (n = 13); OA patients prior to primary arthroplasty (n = 13)	Hyaluronic acid, IL-1B, type III procollagen peptide	
Granchi 1998	case control study/Level III	Aseptic loosening of hip prosthesis (n = 35), healthy controls (n = 25)	IL-1a, IL-1B, IL-6, GM-CSF, IL-2 receptor, TNF-a	Differentiated failed arthroplasties to those with CrCoMo or TiAlV components
Hundri-Haspl 2006	case control study/Level III	Large joint implant loosening (n = 50), large joint arthroplasty without loosening (n = 50), candidates for arthroplasty from advanced OA (n = 50), healthy controls (n = 50)	IL-1B, IL-8, TNF-a	Specific type of large joint prostheses not reported

(Appendix 2) contd.....

Study	Study Design/ Level of Evidence	Comparisons	Outcomes	Other Notes
Streich 2003	case control study/Level III	Aseptic loosening of hip prosthesis (n = 23), fixed hip prosthetic implant (n = 23)	IL-6, GM-CSF, elastase	Significantly more males in healthy donors
Granchi 2006	case control study/Level III	Aseptic loosening of hip implant (n = 36), fixed hip implant (n = 33), severe OA prior to arthroplasty (n = 39), healthy subjects (n = 20)	Osteoprotegerin, RANKL	Significantly more males in healthy donors; significantly longer duration for post-op evaluation in loosening (112 mos) vs. stable joint (32 mos)
Synpniewska 2002	case control study/Level III	aseptic loosening of hip implant (n = 10); primary OA patients undergoing primary arthroplasty (n = 39)	synovial levels: osteocalcin, B-crosslaps, IL-1a, IL-8, IL-10	no fixed arthroplasty control group; excluded patients from analysis if missing outcome data
Kovacik 2000	case control study/Level III	hip implant revisions (n = 20); primary OA patients undergoing primary arthroplasty (n = 7)	synvoial levels: TRAP, IL-1B	does not specific reasons for hip implant revions
Nivbrant 1999	case control study/Level III	loose hip prosthesis (n = 38), fixed hip arthroplasty (n = 12); OA patients undergoing primary arthroplasty (n = 38)	synovial levels: IL-1B, TNF-a, IL-6	mean time from implantation for loose prosthetics was 9.8yrs vs. 2.8yrs in fixed implant; excluded patients from analysis if missing data; median data only
Kawasaki 2001	case control study/Level III	loosened hip prosthesis (n = 5), OA patients from primary hip dysplasia (n = 19), osteonecrosis of femoral neck (n = 21)	synovial levels: YKL-40	average age for OA patients was 48 vs. 63 in loosed implant group; median data only
Nildotter 2007	case control study/Level III	infected hip implants (n = 25), loosened hip implant (n = 60), OA patients undergoing primary arthroplasty (n = 46)	synovial levels: lactate, TNF-a, IL-1B, IL-6 serum levels: CRP, ESR, IL-6, TNF-a, IL-1B	no data provided on serum cytokine levels collected; no comparisons to OA control group provided; excluded patients from analysis if missing data
Bottner 2007	case control study/Level III	septic revision for large joints (n = 21); aseptic revision of large joints (n = 57)	CRP, ESR, IL-6, TNF-a, procalcitonin	Knee/hip revisions: 5/16 for septic, 23/34 for aseptic
Frank 2004	case control study/Level III	staph cultures from prosthetic joint infections compared to coag (-) staph cultures from non-prosthetic joint infections	icaADBC operon	
Galdbert 2000	case control study/Level III	staph epidermidis strains from prosthetic infections (n = 54), staph epidermidis strains from skin flora from 8 healthy individuals	ica operon, biofilm production, genes: clfA, atlE, fbe, fnbA, cna	
Di Cesare 2005	case control study/Level III	58 patients undergoing reoperation of hip or knee: 17 patients with septic prosthetic infection; 41 patients without evidence of infection	CRP, ESR, IL-6	
Rafiq 2000	case control study/Level III	patients with prosthetic infection from gram positive organisms (n = 15), control group of 32 patients – 21 with closed fracture, 11 with primary OA	IgG and IgG to lipoteichoic acid	
Spangehl 1999	prospective cohort study/Level II	prospectively evaluated 178 patients undergoing revision hip replacements	determined specificity, sensitivity for CRP, ESR in prosthetic infections	no specific data provided for comparison analysis; excluded patients from analysis if missing data
Hall 2001	prospective study/Level II	102 patients undergoing primary THA	serum levels: norepinephrine, epinephrine, cortisol, IL-6, CRP; pain, function, and WOMAC ³ at 1 and 6 months;	no control group

(Appendix 2) contd....

Study	Study Design/ Level of Evidence	Comparisons	Outcomes	Other Notes
Wozniak 2004	prospective cohort study/Level II	33 patients undergoing primary TKA or THA due to OA, divided amongst those experiencing pain (n = 8) and no pain (n = 25) after 2-3 years; 14 healthy controls	neutrophils isolated from peripheral blood and analyzed for NO production – at baseline, 2 weeks, 2 months, and 2-3 years post-operatively;	no apparent tool for pain assessment provided; 5/8 reporting pain had evidence of loosening
Messiah 1996	prospective cohort study/Level II	11 patients undergoing revision TKA; ruled out infection;	synovial LDH levels and correlated to presence of polyethylene wear	no control group; no apparent standardized method for measurement of polyethylene wear reported
Wilkinson 2003	case control study/Level III	20 patients undergoing primary THA or OA; compared development of heterotopic ossification by 26 weeks (n = 9) with those without evidence of such (n = 11)	N-terminal propeptide of type-I procollagen, osteocalcin, bone ALP, C-telopeptide of type-I collagen, urine cross-linked N-telopeptide of type-I collagen and free deoxypyridinoline	no control group; only 26 weeks followup for development of heterotopic ossification
Ackland 2007	prospective cohort study/Level II	129 patients undergoing THA or TKA, divided amongst those with hsCRP >3mg/L (n = 63) and those with hsCRP <3mg/L (n = 66)	length of stay, morbid events	

¹RSA = radiostereometric analysis²MTPM = the largest three-dimensional translation of the tibial component³WOMAC = Western Ontario and McMaster universities osteoarthritis index

Level of evidence based on University of Oxford Center for Evidence-based medicine: Level 1: Testing of previously developed diagnostic criteria in series of consecutive patients (with universally applied reference "gold" standard); Systematic review of Level-I studies

Level 2: Development of diagnostic criteria on basis of consecutive patients (with universally applied reference "gold" standard); Systematic review of Level-II studies

Level 3: Study of nonconsecutive patients (without consistently applied reference "gold" standard); Systematic review of Level-III studies

Level 4: Case-control study; Poor reference standard

Level 5: Expert opinion

REFERENCES

- [1] Kurtz S, Ong K, Lau E, Mowat F, Halpern M. Projections of primary and revision hip and knee arthroplasty in the United States from 2005 to 2030. *J Bone Joint Surg Am* 2007; 89: 780-5.
- [2] Biomarkers Definitions Working Group. Biomarkers and surrogate endpoints: Preferred definitions and conceptual framework. *Clin Pharmacol Ther* 2001; 69: 89-95.
- [3] Berlin JA. Does blinding of readers affect the results of meta-analyses? University of Pennsylvania meta-analysis blinding study group. *Lancet* 1997; 350: 185-6.
- [4] Antoniou J, Huk O, Zukor D, et al. Collagen crosslinked n-telopeptides as markers for evaluating particulate osteolysis: A preliminary study. *J Orthop Res* 2000; 18: 64-7.
- [5] von Schewelov T, Carlsson A, Dahlberg L, von Schewelov T, Carlsson A, Dahlberg L. Cross-linked n-telopeptide of type I collagen (ntx) in urine as a predictor of periprosthetic osteolysis. *J Orthop Res* 2006; 24: 1342-8.
- [6] Arabmotlagh M, Sabljic R, Rittmeister M, Arabmotlagh M, Sabljic R, Rittmeister M. Changes of the biochemical markers of bone turnover and periprosthetic bone remodeling after cemented hip arthroplasty. *J Arthroplasty* 2006; 21: 129-34.
- [7] Fiorito S, Magrini L, Goillard C, Fiorito S, Magrini L, Goillard C. Pro-inflammatory and anti-inflammatory circulating cytokines and periprosthetic osteolysis. *J Bone Joint Surg Br* 2003; 85: 1202-6.
- [8] Hernigou P, Intrator L, Bahrami T, Bensussan A, Farset JP. Interleukin-6 in the blood of patients with total hip arthroplasty without loosening. *Clin Orthop Relat Res* 1999; 147-54.
- [9] Schneider U, Breusch SJ, Termaat S, et al. Increased urinary crosslink levels in aseptic loosening of total hip arthroplasty. *J Arthroplasty* 1998; 13: 687-92.
- [10] Li MG, Thorsen K, Nilsson KG, Li MG, Thorsen K, Nilsson KG. Increased bone turnover as reflected by biochemical markers in patients with potentially unstable fixation of the tibial component. *Arch Orthop Trauma Surg* 2004; 124: 404-9.
- [11] Savarino L, Granchi D, Cenni E, et al. Systemic cross-linked n-terminal telopeptide and procollagen I c-terminal extension peptide as markers of bone turnover after total hip arthroplasty. *J Bone Joint Surg Br* 2005; 87: 571-6.
- [12] Wilkinson JM, Hamer AJ, Rogers A, et al. Bone mineral density and biochemical markers of bone turnover in aseptic loosening after total hip arthroplasty. *J Orthop Res* 2003; 21: 691-6.
- [13] Moreschini O, Fiorito S, Magrini L, et al. Markers of connective tissue activation in aseptic hip prosthetic loosening. *J Arthroplasty* 1997; 12: 695-703.
- [14] Granchi D, Verri E, Ciapetti G, et al. Bone-resorbing cytokines in serum of patients with aseptic loosening of hip prostheses. *J Bone Joint Surg Br* 1998; 80: 912-7.
- [15] Hundrie-Haspl Z, Pecina M, Haspl M, et al. Plasma cytokines as markers of aseptic prosthesis loosening. *Clin Orthop* 2006; 453: 299-304.
- [16] Streich NA, Breusch SJ, Schneider U, Streich NA, Breusch SJ, Schneider U. Serum levels of interleukin 6 (il-6), granulocyte-macrophage colony-stimulating factor (gm-csf) and elastase in aseptic prosthetic loosening. *Int Orthop* 2003; 27: 267-71.
- [17] Granchi D, Pellacani A, Spina M, et al. Serum levels of osteoprotegerin and receptor activator of nuclear factor-kappaB ligand as markers of periprosthetic osteolysis. *J Bone Joint Surg Am* 2006; 88: 1501-9.
- [18] Nivbrant B, Karlsson K, Karrholm J. Cytokine levels in synovial fluid from hips with well-functioning or loose prostheses. *J Bone Joint Surg Br* 1999; 81: 163-6.
- [19] Kovacik MW, Gradisar IA, Jr., Haprian JJ, et al. Osteolytic indicators found in total knee arthroplasty synovial fluid aspirates. *Clin Orthop* 2000; 186-94.
- [20] Kawasaki M, Hasegawa Y, Kondo S, et al. Concentration and localization of ykl-40 in hip joint diseases. *J Rheumatol* 2001; 28: 341-5.
- [21] Sypniewska G, Lis K, Bilinski PJ, Sypniewska G, Lis K, Bilinski PJ. Bone turnover markers and cytokines in joint fluid: Analyses in

- 10 patients with loose hip prosthesis and 39 with coxarthrosis. *Acta Orthop Scand* 2002; 73: 518-22.
- [22] Rafiq M, Worthington T, Tebbs SE, et al. Serological detection of gram-positive bacterial infection around prostheses. *J Bone Joint Surg Br* 2000; 82: 1156-61.
- [23] Galdbart JO, Allignet J, Tung HS, et al. Screening for staphylococcus epidermidis markers discriminating between skin-flora strains and those responsible for infections of joint prostheses. *J Infect Dis* 2000; 182: 351-5.
- [24] Frank KL, Hanssen AD, Patel R, Frank KL, Hanssen AD, Patel R. Icaaa is not a useful diagnostic marker for prosthetic joint infection. *J Clin Microbiol* 2004; 42: 4846-9.
- [25] Di Cesare PE, Chang E, Preston CF, et al. Serum interleukin-6 as a marker of periprosthetic infection following total hip and knee arthroplasty. *J Bone Joint Surg Am* 2005; 87: 1921-7.
- [26] Bottner F, Wegner A, Winkelmann W, et al. Interleukin-6, procalcitonin and tnf-alpha: Markers of peri-prosthetic infection following total joint replacement. *J Bone Joint Surg Br* 2007; 89: 94-9.
- [27] Nilsdotter-Augustinsson A, Briheim G, Herder A, et al. Inflammatory response in 85 patients with loosened hip prostheses: A prospective study comparing inflammatory markers in patients with aseptic and septic prosthetic loosening. *Acta Orthop* 2007; 78: 629-39.
- [28] Spangehl MJ, Masri BA, O'Connell JX, Duncan CP. Prospective analysis of preoperative and intraoperative investigations for the diagnosis of infection at the sites of two hundred and two revision total hip arthroplasties. *J Bone Joint Surg Am* 1999; 81: 672-83.
- [29] Hall GM, Peerbhay D, Shenkin A, et al. Relationship of the functional recovery after hip arthroplasty to the neuroendocrine and inflammatory responses. [see comment]. *Br J Anaesth* 2001; 87: 537-42.
- [30] Wozniak W, Markuszewski J, Wierusz-Kozlowska M, et al. Neutrophils are active in total joint implant loosening. *Acta Orthop Scand* 2004; 75: 549-53.
- [31] Wilkinson JM, Stockley I, Hamer AJ, et al. Biochemical markers of bone turnover and development of heterotopic ossification after total hip arthroplasty. *J Orthop Res* 2003; 21: 529-34.
- [32] Messieh M, Messieh M. Synovial fluid levels of lactate dehydrogenase in patients with total knee arthroplasty. *J Arthroplasty* 1996; 11: 484-6.
- [33] Ackland GL, Scollay JM, Parks RW, et al. Pre-operative high sensitivity c-reactive protein and postoperative outcome in patients undergoing elective orthopaedic surgery. *Anaesthesia* 2007; 62: 888-94.
- [34] Bozic KJ, Kurtz SM, Lau E, Ong K, Vail TP, Berry DJ. The epidemiology of revision total hip arthroplasty in the united states. *J Bone Joint Surg Am* 2009; 91: 128-33.
- [35] Perry MJ, Ponsford FM, Mortua FY, Learmonth ID, Atkins RM, Elson CJ. Osteolytic properties of the synovial-like tissue from aseptically failed joint prostheses. *Br J Rheumatol* 1996; 35: 943-50.
- [36] Xu JW, Li TF, Partsch G, Ceponis A, Santavirta S, Konttinen YT. Interleukin-11 (il-11) in aseptic loosening of total hip replacement (thr). *Scand J Rheumatol* 1998; 27: 363-7.
- [37] Lassus J, Waris V, Xu JW, et al. Increased interleukin-8 (il-8) expression is related to aseptic loosening of total hip replacement. *Arch Orthop Trauma Surg* 2000; 120: 328-32.
- [38] Waddell J, Pritzker KP, Boynton EL. Increased cytokine secretion in patients with failed implants compared with patients with primary implants. *Clin Orthop Relat Res* 2005; 434: 170-6.

Received: October 9, 2009

Revised: April 4, 2010

Accepted: July 7, 2010

© Mertens and Singh; Licensee Bentham Open.

This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.